



Seasonal dynamics of soil water content in the typical vegetation and its response to precipitation in a semi-arid area of Chinese Loess Plateau

ZHOU Tairan^{1,2}, HAN Chun^{1,2}, QIAO Linjie^{1,2}, REN Chaojie^{1,2}, WEN Tao^{1,2},
ZHAO Changming^{1,2*}

¹ State Key Laboratory of Grassland and Agro-Ecosystems, School of Life Sciences, Lanzhou University, Lanzhou 730000, China;

² Gansu Provincial Field Scientific Observation and Research Station of Mountain Ecosystems, Lanzhou University, Lanzhou 730000, China

Abstract: Soil water content is a key limiting factor for vegetation growth in the semi-arid area of Chinese Loess Plateau and precipitation is the main source of soil water content in this area. To further understand the impact of vegetation types and environmental factors such as precipitation on soil water content, we continuously monitored the seasonal dynamics in soil water content in four plots (natural grassland, *Caragana korshinskii*, *Armeniaca sibirica* and *Pinus tabulaeformis*) in Chinese Loess Plateau. The results show that the amplitude of soil water content fluctuation decreases with an increase in soil depth, showing obvious seasonal variations. Soil water content of artificial vegetation was found to be significantly lower than that of natural grassland, and most precipitation events have difficulty replenishing soil water content below a depth of 40 cm. Spring and autumn are the key seasons for replenishment of soil water by precipitation. Changes in soil water content are affected by precipitation, vegetation types, soil evaporation and other factors. The interception effect of vegetation on precipitation and the demand for water consumption by transpiration are the key factors affecting the efficiency of soil water replenishment by precipitation in this area. Due to artificial vegetation plantation in this area, soil will face a water deficit crisis in the future.

Keywords: soil water content; vegetation type; precipitation; seasonal change; evaporation

Citation: ZHOU Tairan, HAN Chun, QIAO Linjie, REN Chaojie, WEN Tao, ZHAO Changming. 2021. Seasonal dynamics of soil water content in the typical vegetation and its response to precipitation in a semi-arid area of Chinese Loess Plateau. Journal of Arid Land, 13(10): 1015–1025. <https://doi.org/10.1007/s40333-021-0021-5>

1 Introduction

Soil water content is a key factor for the growth and survival of vegetation in semi-arid areas (Shao et al., 2014), and plays an important role in the ecohydrological cycle of ecosystem (Daly and Porporato, 2005). At the plot scale, the independent and interactive effects of precipitation, vegetation types, soil and other factors make soil water content extremely heterogeneous (Grayson et al., 1997). Precipitation is the main source of soil water content replenishment in semi-arid areas, as well as a key factor affecting soil water content (Wilson et al., 2004). Due to the high potential evapotranspiration in semi-arid areas, small precipitation events with less amount

*Corresponding author: ZHAO Changming (E-mail: zhaochm@lzu.edu.cn)

Received 2021-06-30; revised 2021-10-18; accepted 2021-10-21

© Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences, Science Press and Springer-Verlag GmbH Germany, part of Springer Nature 2021

often do not have a significant impact on soil water content (Kurc and Small, 2007). When precipitation amount increases, water can penetrate deeper soil profiles (Schwinning and Sala, 2004). Additionally, vegetation type plays a key role in the effectiveness of precipitation (Schlueter et al., 2013). It can affect the process of soil water input through canopy precipitation interception, hydraulic enhancement and soil organic matter content (Odindi and Kakembo, 2011), and can affect the output of soil water through processes such as evaporation, transpiration and infiltration (Pan et al., 2008). Vegetation can regulate soil water input and output, reaching a stable dynamic equilibrium state (Yu et al., 2018), and resulting in continuous depletion of soil water (Zhao et al., 2017). For instance, soil water depletion by transpiration in shrubs tends to be more severe than that in grasslands (Schlueter et al., 2013). Therefore, vegetation type is a key regulating factor of water cycle in semi-arid areas where water resources are scarce (Wu et al., 2016). The patterns of change in soil water content under different vegetation coverage, depth and season were significantly different (Yu et al., 2017; Hou et al., 2018).

The Loess Plateau in China is the largest loess accumulation area in the world (D'Odorico et al., 2007), and has long faced many ecological problems such as serious soil erosion and vegetation degradation (Zhang et al., 2008). To control soil erosion and improve the environment, China has implemented the world's largest "Grain for Green" project in this area (Zhou et al., 2012). The "Grain for Green" project has significantly increased the vegetation coverage and carbon sequestration potential in the semi-arid areas of western China (Zhou et al., 2012). Nonetheless, it also increased soil water consumption (Zhang et al., 2020), thus leading to a series of ecological and environmental problems such as deep soil drying in forest lands (Liang et al., 2018). In the water-deficient environment of the semi-arid areas of the Loess Plateau, different vegetation types have adopted different water use strategies (Hou et al., 2018; Yu et al., 2018), and these influences on hydrological processes such as soil water content storage, migration and transformation are quite different (Cheng et al., 2020).

In recent years, global warming has exacerbated the problem of soil water shortage in the semi-arid areas (Zhang et al., 2020), and the factors affecting soil water content may change drastically with the seasons (Zhu et al., 2014). Seasonal dynamics of soil water content under different vegetation types in the semi-arid areas have become an important research topic in ecohydrology (Mei et al., 2019; Yu et al., 2019). It is necessary to further quantify the influence of precipitation and vegetation types and other factors on the heterogeneity of soil water content in the semi-arid areas (Zhao et al., 2017; Yu et al., 2018). Therefore, we used TEROS12 sensors (METER Group, Pullman, USA) to monitor the soil water content of the typical vegetation in a semi-arid area of Chinese Loess Plateau between 1 January and 31 December, 2020. For real-time monitoring of meteorological factors, the dynamics of soil water content in different soil depths and the differences in seasonal response patterns to precipitation of four vegetation types (natural grassland, *Caragana korshinskii*, *Pinus tabulaeformis* and *Armeniaca sibirica*) were compared.

2 Materials and methods

2.1 Study area

This study was conducted at the Gansu Provincial Field Scientific Observation and Research Station of Mountain Ecosystems, China (35°58'20"N, 104°20'6"E). The region has an average altitude of 2000 m a.s.l, annual average temperature of 7°C and average annual precipitation of 300–350 mm. We selected sample plots with the typical vegetation for 60 a plantation to monitor soil water content. Four vegetation types, i.e., natural grassland, *C. korshinskii*, *A. sibirica* and *P. tabulaeformis* were selected with similar environmental conditions and planting ages (Fig. 1). The main vegetation in natural grassland included *Artemisa sacrorum*, *Ajania fruticulosa* and *Stipa bungeana*. The understory vegetation types of other three plots were dominated by *A. sacrorum*. The vegetation types, total vegetation coverage, soil bulk density and soil particle size composition of the sample plots are shown in Table 1.




Fig. 1 Landscape of natural grassland (a), *Caragana korshinskii* (b), *Armeniaca sibirica* (c) and *Pinus tabulaeformis* (d)

Table 1 Information of vegetation and soil of the sample plot

Index	Natural grassland	<i>Caragana korshinskii</i>	<i>Armeniaca sibirica</i>	<i>Pinus tabulaeformis</i>
Stand age (a)	-	56	50	50
Altitude (m)	2050	2084	2156	2120
Slope (°)	15	20	15	18
Aspect	Southeast	Southeast	Southeast	West
Total vegetation coverage (%)	95	57	50	75
Dominant species	<i>A. sacrorum</i> , <i>Ajania fruticulosa</i> , <i>Stipa bungeana</i>	<i>A. sacrorum</i> , <i>Caragana jubata</i> , <i>Achnatherum splendens</i>	<i>A. sacrorum</i> , <i>Ajania fruticulosa</i> , <i>Stipa capillata</i>	<i>A. sacrorum</i> , <i>Stipa bungeana</i> , <i>Aster altaicus</i>
Understory vegetation coverage (%)	-	22	25	15
Bulk density (g/cm ³)	1.19±0.05	0.98±0.07	1.07±0.04	1.08±0.05
Clay (%)	8.36±1.68	7.77±0.37	6.37±0.94	6.86±0.89
Silt (%)	81.25±4.28	78.83±1.63	74.61±0.5	75.69±0.83
Sand (%)	10.39±2.69	13.39±1.64	19.01±1.07	17.45±1.66

Note: - means no value. Mean±SE.

2.2 Experimental design and data collection

The TEROS12 soil moisture sensor (METER Group, Inc., USA) were installed at 10, 20, 40, 60, and 80 cm depths with an interval of 10 min to measure soil water content. We calibrated soil water content data according to the empirical calibration equation to obtain an accuracy of approximately 1%–2%. Meteorological data such as precipitation, air temperature and relative humidity were measured at 10 min intervals using the automatic monitoring probe of Dynamet's scientific research-level automatic weather station, and the vapor pressure deficit was fitted using an empirical formula. The accuracy of precipitation data was 4%, as recorded on the CR1000 data collector (CR1000, Campbell Scientific, Logan, Utah, USA).

2.3 Data analysis

We used one-way analysis of variance (ANOVA) and least significant difference method (LSD) to analyze the differences in soil water content between different soil depth in each plot. Linear regression analysis was used to obtain the response model of soil water storage (SWS) to precipitation. SWS capacity was calculated as follows:

$$\text{SWS}_{0-90} = \theta_{10} \times 15 + \theta_{20} \times 15 + \theta_{40} \times 20 + \theta_{60} \times 20 + \theta_{80} \times 20, \quad (1)$$

where SWS_{0-90} is the soil water storage capacity of 0–90 cm soil depth (cm); and θ_{10} , θ_{20} , θ_{40} , θ_{60} and θ_{80} are the soil water content at depths of 10, 20, 40, 60 and 80 cm, respectively (cm) (Mei et al., 2019). The change in SWS capacity (ΔSWS , cm) was calculated as follows:

$$\Delta\text{SWS} = \text{SWS}_2 - \text{SWS}_1, \quad (2)$$

where SWS_1 is the soil water storage before a precipitation event (cm); and SWS_2 is the peak

soil water storage following a precipitation event (cm). ΔSWS is used to characterize the supply capacity of precipitation to soil water. Coefficient of variation (CV) is calculated as follows:

$$\text{CV} = \sigma / \mu \times 100\%, \quad (3)$$

where μ is the average value of soil sample; and σ is the standard deviation of sample (Xu et al., 2017). All statistical analyses were performed using SPSS software (version 20.0; SPSS Inc., Chicago, IL, USA). Graphics were drawn using Origin 8.5 (Origin Lab, Northampton, MA, USA).

3 Results

3.1 Precipitation

Figure 2 shows that 73 precipitation events occurred during the study period, with summer accounting for 62.73% of annual precipitation (338.3 mm). Precipitation events below 5 mm were the most frequent, but the total amount accounted for only 13.09% of annual precipitation. Precipitation events of 5–10 mm and 10–20 mm occurred in spring, summer and autumn and were mainly concentrated in summer, accounting for 23.00% and 42.21% of annual precipitation, respectively. Precipitation events greater than 20 mm only occurred in summer, accounting for 34.59% of summer precipitation and 21.70% of annual precipitation.




Fig. 2 Frequency (a) and amount (b) of precipitation in each season

3.2 Soil water content

Table 2 shows variation in average soil water content of each soil depth for different vegetation types. The average soil water content exhibited the following trend: natural grassland > *P. tabulaeformis* > *C. korshinskii* > *A. sibirica*; CV associated with soil water content exhibited the following sequence: *P. tabulaeformis* > *C. korshinskii* > *A. sibirica* > natural grassland. These results indicated that soil water content of natural grassland exhibited a small variation. The changes in the daily average temperature and vapor pressure deficit reflect the typical continental monsoon climate in the area, with rapid changes being observed in spring and autumn. The climatic characteristics of rain and heat drastically altered soil water content (Fig. 3). The analysis of variance showed that the difference between the maximum and minimum values of soil water content of each soil depth was significant ($P < 0.05$), indicating that soil water content was significantly affected by vegetation, precipitation and other environmental factors.

Soil water content of different vegetation types varied with soil depth. The maximum soil water content appeared in 10 cm soil depth; however, soil water content in 10 cm soil depth was significantly lower than that in 20 cm soil depth ($P < 0.05$). Except for *C. korshinskii* plot, CV in soil water content in 10 cm soil depth was the largest, and the minimum soil water content was observed in 80 cm soil depth, which indicated that precipitation had the highest effect on 10 cm soil depth. Soil water content in 40 cm soil depth of *P. tabulaeformis* plot was higher than those in other soil depths, while the maximum soil water content appeared in 20 cm soil depth, indicating that this soil layer is important for water storage.

Table 2 Statistical values of soil water content in the four plots

Sample plot	Soil depth (cm)	Average (%)	Maximum (%)	Minimum (%)	SD (%)	CV (%)
Natural grassland (<i>n</i> =52,704)	10	21.38 ^c	34.24	12.40	3.66	17.13
	20	23.93 ^a	33.47	17.50	2.75	11.51
	40	20.12 ^c	25.62	14.06	2.96	14.70
	60	17.61 ^e	20.20	11.72	2.45	13.93
	80	14.20 ^g	17.02	10.31	2.00	14.07
<i>C. korshinskii</i> (<i>n</i> =52,704)	10	18.53 ^d	33.51	11.79	4.20	22.64
	20	20.87 ^c	30.22	12.61	4.79	22.95
	40	9.87 ^j	15.21	4.93	2.98	30.17
	60	9.25 ^k	12.76	6.42	1.73	18.70
	80	8.27 ^l	9.70	6.93	0.47	5.74
<i>A. sibirica</i> (<i>n</i> =52,704)	10	17.40 ^e	30.66	10.11	4.78	27.45
	20	18.57 ^d	27.32	12.53	3.53	19.00
	40	12.69 ⁱ	18.21	8.12	2.57	20.27
	60	4.25 ⁿ	4.94	3.03	0.42	9.96
	80	4.79 ^m	6.41	3.03	0.96	20.15
<i>P. tabulaeformis</i> (<i>n</i> =52,704)	10	13.21 ^h	30.02	5.64	4.73	35.82
	20	20.24 ^c	37.49	11.71	4.05	19.99
	40	23.01 ^b	31.71	14.72	2.90	12.62
	60	16.34 ^f	19.28	10.58	2.60	15.88
	80	8.15 ^l	11.52	3.85	2.30	28.26

Note: SD, standard deviation; CV, coefficient of variation; Different lowercase letters within the same column showed significant difference at $P < 0.05$ level.

The results in Figure 3 also show fluctuations in soil water content of each soil depth during experimental period. The peak value of 10 cm soil depth corresponded strongly with precipitation event, whereas soil water content fluctuations in deeper depth were observed with the large and continuous precipitation event in summer. Otherwise, soil water content remained relatively stable during experimental period. In summer, when precipitation was relatively concentrated, soil water content in deeper depths (60 and 80 cm) dropped to the lowest (Fig. 3d and e). Except for natural grassland, the maximum water content in 80 cm soil depth of other plots was less than 12%, indicating that soil water content had been depleted during experimental period. Except for natural grassland, soil water content became low at the end of experiment, and there was a negative balance of soil water content during experimental period (Fig. 3).

3.3 Response of soil water content to precipitation

Precipitation events of less than 5 mm had little impact on soil water content. We selected three precipitation events (5–10, 10–20 and >20 mm) and evaluated their effects on soil water content (Fig. 4). The response patterns of each plot to 6.8 mm precipitation were similar (Fig. 4a1, b1, c1 and d1). Precipitation of 6.8 mm increased soil water content in a depth of 10 cm, and infiltrated to 20 cm soil depth after 1 d of precipitation. The effect of 18.4 mm precipitation on soil water content of natural grassland and *C. korshinskii* plot was similar to that of 6.8 mm precipitation (Fig. 4a2, b2, c2 and d2). Soil water content of *A. sibirica* and *P. tabulaeformis* plots penetrated into 40 cm soil depth after 2 d of 18.4 mm precipitation (Fig. 4c3 and d3). Precipitation of 25.1 mm caused soil water content in natural grassland and *P. tabulaeformis* plots to infiltrate the depth of 80 cm (Fig. 4a3 and c3), and *C. korshinskii* and *A. sibirica* plots infiltrated to the depth of 40–60 cm (Fig. 4b3 and d3). With the increase in precipitation, soil water content varied at different soil depths (Fig. 4). Natural grassland and *P. tabulaeformis* plot showed the most variation of soil water content. The 25.1 mm precipitation was the largest event that occurred during study period.




Fig. 3 Daily time series of precipitation (a), air temperature (f), vapor pressure deficit (f) and soil water content in natural grassland, *C. korshinskii*, *A. sibirica*, and *P. tabulaeformis* plots at soil depths of 10 (a), 20 (b), 40 (c), 60 (d) and 80 cm (e)

The magnitude of this precipitation event was sufficient to replenish soil water content to the depth of 80 cm in natural grassland and *P. tabulaeformis* plot after 3 d of precipitation.

3.4 Dynamics of SWS capacity and response to precipitation

SWS capacity of natural grassland was significantly higher than those of other plots ($P < 0.05$), and that of *A. sibirica* was the lowest ($P < 0.05$; Table 3). Except for natural grassland in spring, SWS capacity was significantly higher in autumn than in other seasons ($P < 0.05$), and had a low CV (Table 3). Annual SWS capacity reached its lowest value in summer. Annual SWS exhibited the following descending order: natural grassland > *P. tabulaeformis* > *C. korshinskii* > *A. sibirica*. Natural grassland exhibited the lowest CV in SWS capacity, while *C. korshinskii* was the highest.

SWS and precipitation were significantly correlated during experimental period ($P < 0.05$). Seasonal variation of SWS capacity is obvious (Fig. 5). Except in winter, response of SWS to precipitation in *P. tabulaeformis* plot was higher than those of other plots (Fig. 5a, b and c); however, in winter, the response of SWS capacity in natural grassland was the strongest (Fig. 5d). The efficiency of SWS capacity was the highest in spring for *A. sibirica* and *P. tabulaeformis* plots (Fig. 5a), followed by those in winter (Fig. 5d) and summer (Fig. 5b). For natural grassland

and *C. korshinskii* plot, the highest SWS capacity occurred in winter (Fig. 5d), followed by spring (Fig. 5a) and autumn (Fig. 5c).




Fig. 4 Response of soil water content to precipitation events (6.8, 18.4 and 25.1 mm) in different soil depths of the four plots. (a1–a3), natural grassland; (b1–b3), *C. korshinskii*; (c1–c3), *A. sibirica*; (d1–d3), *P. tabulaeformis*. P, precipitation.

4 Discussion


4.1 Variation in soil water content and its influencing factors

The results of the study indicate that the trend of soil water content in each vegetation type is consistent with the change in precipitation (Fig. 3), indicating that precipitation is an important factor affecting soil water content. However, the response of soil water content to precipitation is more complicated, and factors such as vegetation type, soil depth and environmental factors affect the dynamics of soil water content. Soil evaporation and vegetation transpiration patterns in different types of vegetation are main reasons for the dynamics of soil water content (Kakembo, 2009). Our results confirmed that annual SWS capacity for different types of vegetation varied significantly (Table 3), and SWS capacity of natural grassland was much higher than those of other plots. Soil water content of *A. sibirica* plot had dropped owing to the depleting of water content (Laio et al., 2001), which is undeniably dangerous for vegetation. Low soil water content for a prolonged period limits the growth of vegetation and eventually causes vegetation death (Yu

Table 3 Statistical value of soil water storage capacity during experimental period

Season	Plot	Average (cm)	SD (cm)	CV (%)
Spring (n=92)	Natural grassland	18.20 ^a	0.81	4.43
	<i>C. korshinskii</i>	11.67 ^e	1.08	9.24
	<i>A. sibirica</i>	9.34 ^k	1.02	10.89
	<i>P. tabulaeformis</i>	15.49 ^c	1.02	6.60
Summer (n=92)	Natural grassland	15.41 ^c	1.95	12.62
	<i>C. korshinskii</i>	10.01 ⁱ	1.88	18.80
	<i>A. sibirica</i>	8.72 ^k	1.91	21.93
	<i>P. tabulaeformis</i>	13.00 ^e	2.67	20.55
Autumn (n=91)	Natural grassland	18.38 ^a	0.40	2.18
	<i>C. korshinskii</i>	13.40 ^d	0.36	2.71
	<i>A. sibirica</i>	11.36 ^g	0.56	4.96
	<i>P. tabulaeformis</i>	15.68 ^c	0.77	4.91
Winter (n=91)	Natural grassland	16.75 ^c	0.89	5.29
	<i>C. korshinskii</i>	10.49 ^h	1.42	13.50
	<i>A. sibirica</i>	9.57 ^j	0.82	8.55
	<i>P. tabulaeformis</i>	13.91 ^c	0.81	5.82
Annual (n=366)	Natural grassland	17.18 ^b	1.67	9.72
	<i>C. korshinskii</i>	11.39 ^f	1.85	16.21
	<i>A. sibirica</i>	9.74 ^j	1.54	15.82
	<i>P. tabulaeformis</i>	14.52 ^c	1.90	13.07

Note: SD, standard deviation; CV, coefficient of variation. Different lowercase letters within the same column showed significant difference among different seasons at $P < 0.05$ level.

**Fig. 5** Response model of change of soil water storage to precipitation

et al., 2018; Yu et al., 2019).

Soil water content decreases with increasing soil depth owing to the weak precipitation influence to deeper soil depth (Fu et al., 2003; Zhao et al., 2017). As shown in Figure 3a, the surface soil water content in 10 cm depth was most sensitive to precipitation with the highest value. As soil depth increased, soil structure became more compact, reducing water infiltration capacity, thus stabilizing deep soil water content. Our result found that 60–80 cm soil depth had the lowest soil water content. Furthermore, CV value in soil water content reflected its stability. The lower the value of soil water content in deep soil, the lesser it was affected by precipitation and other factors (Yu et al., 2017). Vegetation root in deep soil has become an important factor affecting soil water content (Fu et al., 2003). *C. korshinskii*, *A. sibirica* and *P. tabulaeformis* plots all showed the larger CVs in 40–80 cm soil depth (Table 2) because artificial vegetation has a stronger root system at this depth (Xu et al., 2017).

Results showed that soil water content of different types of vegetation displayed seasonal differences (Fig. 3f) due to large differences in factors such as surface evapotranspiration and vegetation transpiration in different seasons (Li et al., 2013; Liu et al., 2017). Soil water content is relatively high in spring, but it also begins to deplete rapidly. This is because, temperature and vapor pressure deficit, vegetation transpiration and soil evaporation increases rapidly (Laio et al., 2001). Both temperature and vapor pressure deficit are relatively high in summer, with obvious fluctuation of rain and heat. The relatively strong input and output processes of soil water content in summer can also be reflected by the higher CV in SWS capacity. Therefore, although there is relatively high frequency of precipitation in summer, SWS capacity does not get supplement significantly. In autumn, temperature and vapor pressure deficit reduce significantly, effects of vegetation transpiration, water consumption and soil evaporation are significantly weakened, and the change in soil water content tends to stabilize (Kakembo, 2009). Autumn is the main supplementary season for soil water content. Additionally, average SWS capacity of each plot is relatively high in autumn.

4.2 Response of soil water content to precipitation

Results of the study indicate that a precipitation of more than 20 mm, which only occurs in summer, is the only way to replenish moisture in deeper soil (>40 cm) (Fig. 4). Precipitation (>20 mm) is important for sustainable growth and survival of vegetation (Li et al., 2013). Precipitation event of 5–20 mm in spring and autumn may be the main source of soil water replenishment in this area (Wang et al., 2013; Mei et al., 2019). In general, most of small-scale precipitation event in this area is quickly offset by evaporation because of limited infiltration depth, whereas large-scale precipitation event would increase infiltration depth and correspondingly reduce evaporation of surface soil water (Mei et al., 2019). Additionally, continuous small-scale precipitation events may have a superimposing effect, i.e., increasing the efficiency of soil water replenishment by increasing surface soil water content, the importance of which cannot be neglected (Peel, 2009). However, we did not observe a significant precipitation superimposing effect during experimental period. This may be due to continuous precipitation event that mainly occurred in summer (Fig. 3), and high temperature and evapotranspiration offset the continuous precipitation event (Laio et al., 2001). Response of soil water content to precipitation has seasonal differences, and the impact and supplement of precipitation in spring and autumn are more obvious due to canopy interception, vegetation transpiration and reduced soil evaporation loss (He and Huang, 2001).

P. tabulaeformis has a higher conversion efficiency of SWS capacity for precipitation due to the combined effect of several factors. Firstly, differences in soil physical and chemical properties are important factors that affect the conversion of precipitation into soil water content. Soil organic matter content and soil porosity of *P. tabulaeformis* plot are relatively high, and soil infiltration rate was higher than those of other plots (Fig. 4) (Guo et al., 2019). Soil water content in 40 cm depth of *C. korshinskii* and *A. sibirica* plots reached the level of dry soil layer of Loess Plateau, and average soil water content reached the level of severe dry layer (Liang et al., 2018). This indicates that the roots of *C. korshinskii* and *P. apricot* plots have a strong ability to absorb and

consume excessive soil water, which makes precipitation difficult to infiltrate to this depth and form a moist layer (Ma et al., 2017). Secondly, research has found that *P. tabulaeformis* adopts active drought resistance strategies such as rainfall interception (Ma et al., 2019) to reduce initial water loss due to transpiration (Liu et al., 2017; Ma et al., 2017). Therefore, *P. tabulaeformis* had a higher efficiency of SWS capacity (Fig. 5).

Measures for vegetation restoration have been carried out for approximately 60 a, and irrigation or other sources of water replenishment except for precipitation in this study area are not existed. Thus, deep water in deep soil depth in artificial plantation forest obtained from precipitation is limited (Fig. 4). Therefore, in the context of future climate change, an increase in the extreme drought event may cause artificial vegetation to face serious soil water deficit crisis (An et al., 2017). Therefore, afforestation in the future should be more cautious. The optimal season for afforestation is spring (Zhou et al., 2012; Ma et al., 2019) due to the high soil water replenishment efficiency (Liu et al., 2017). At the same time, it is essential to implement necessary management and irrigation measures for existing artificial vegetation (Gao et al., 2015) for the demand of soil water content of deep soil depth (Zhao et al., 2017; Mei et al., 2019).

5 Conclusions

Our study confirmed that variation of soil water content in the semi-arid area of Chinese Loess Plateau is significantly affected by precipitation and vegetation type. Moderate precipitation event is the main water source for surface soil. Due to the little effect of precipitation on the deeper soil depth, especially in summer for its high temperature and evaporation, artificial vegetation in this area still faces the water deficit crisis. The order of SWC of vegetation is natural grassland > *P. tabulaeformis* > *C. korshinskii* > *A. sibirica*. Thus, natural grassland is more suitable for the restoration of this area. Therefore, proportion of artificial vegetation and natural restoration should be optimized for sustainable development of water resources in Chinese Loess Plateau.

Acknowledgements

This work was supported by the Strategic Priority Research Program of Chinese Academy of Sciences (XDA20100101), the Major Special Science and Technology Project of Gansu Province, China (18ZD2FA009), and the National Natural Science Foundation of China (31522013).

References

- An W, Li Z, Wang S, et al. 2017. Exploring the effects of the "Grain for Green" program on the differences in soil water in the semi-arid Loess Plateau of China. *Ecological Engineering*, 107: 144–151.
- Cheng R R, Chen Q W, Zhang J G, et al. 2020. Soil moisture variations in response to precipitation in different vegetation types: A multi-year study in the loess hilly region in China. *Ecohydrology*, 13(3): e2196, doi: 10.1002/eco.2196.
- Daly E, Porporato A. 2005. A review of soil moisture dynamics: From rainfall infiltration to ecosystem response. *Environmental Engineering Science*, 22(1): 9–24.
- D'Odorico P, Caylor K, Okin G S, et al. 2007. On soil moisture–vegetation feedbacks and their possible effects on the dynamics of dryland ecosystems. *Journal of Geophysical Research*, 112: G04010, doi: 10.1029/2006JG000379.
- Fu B J, Wang J, Chen L D, et al. 2003. The effects of land use on soil moisture variation in the Danangou catchment of the Loess Plateau, China. *CATENA*, 54(1–2): 197–213.
- Gao L, Shao M G, Peng X H, et al. 2015. Spatio-temporal variability and temporal stability of water contents distributed within soil profiles at a hillslope scale. *CATENA*, 132: 29–36.
- Grayson R B, Western A W, Chiew F H S, et al. 1997. Preferred states in spatial soil moisture patterns: Local and nonlocal controls. *Water Resources Research*, 33(12): 2897–2908.
- Guo S J, Xu Y D, He C, et al. 2019. Differential responses of soil quality in revegetation types to precipitation gradients on the Loess Plateau. *Agricultural and Forest Meteorology*, 276–277: 107622, doi: 10.1016/j.agrformet.2019.107622.
- He X B, Huang Z B. 2001. Zeolite application for enhancing water infiltration and retention in loess soil. *Resources Conservation and Recycling*, 34(1): 45–52.
- Hou G, Bi H, Wei X, et al. 2018. Response of soil moisture to single rainfall events under three vegetation types in the Gully Region of the Loess Plateau. *Sustainability*, 10(10): 3793.

- Kakembo V. 2009. Vegetation patchiness and implications for landscape function: The case of *Pteronia incana* invader species in Ngqushwa Rural Municipality, Eastern Cape, South Africa. *CATENA*, 77(3): 180–186.
- Kure S A, Small E E. 2007. Soil moisture variations and ecosystem–scale fluxes of water and carbon in semiarid grassland and shrubland. *Water Resources Research*, 43(6): W06416, doi: 10.1029/2006WR005011.
- Liao F, Porporato A, Ridolfi L, et al. 2001. Plants in water-controlled ecosystems: active role in hydrologic processes and response to water stress II. Probabilistic soil moisture dynamics. *Advances in Water Resources*, 24(7): 707–723.
- Li X Y, Zhang S Y, Peng H Y, et al. 2013. Soil water and temperature dynamics in shrub–encroached grasslands and climatic implications: Results from Inner Mongolia steppe ecosystem of North China. *Agricultural and Forest Meteorology*, 171–172: 20–30.
- Liang H, Xue Y, Li Z, et al. 2018. Soil moisture decline following the plantation of *Robinia pseudoacacia* forests: Evidence from the Loess Plateau. *Forest Ecology and Management*, 412: 62–69.
- Liu X, Zhang B, Zhuang J Y, et al. 2017. The relationship between sap flow density and environmental factors in the Yangtze River Delta Region of China. *Forests*, 8(3): 74.
- Ma C, Luo Y, Shao M, et al. 2017. Environmental controls on sap flow in black locust forest in Loess Plateau, China. *Scientific Reports*, 7(1): 13160, doi: 10.1038/s41598-017-13532-8.
- Ma C K, Li X D, Luo Y, et al. 2019. The modelling of rainfall interception in growing and dormant seasons for a pine plantation and a black locust plantation in semi-arid Northwest China. *Journal of Hydrology*, 577: 123849, doi: 10.1016/j.jhydrol.2019.06.021.
- Mei X M, Ma L, Zhu Q K, et al. 2019. The variability in soil water storage on the loess hillslopes in China and its estimation. *Catena*, 172: 807–818.
- Ondindi J O, Kakembo V. 2011. The hydrological response of *Pteronia incana*–invaded areas in the Eastern Cape Province, South Africa. *Ecohydrology*, 4(6): 832–840.
- Pan M, Wood E F, Wojcik R, et al. 2008. Estimation of regional terrestrial water cycle using multi–sensor remote sensing observations and data assimilation. *Remote Sensing of Environment*, 112(4): 1282–1294.
- Peel M C. 2009. Hydrology: catchment vegetation and runoff. *Progress in Physical Geography*, 33(6): 837–844.
- Schlueter S, Vogel H J, Ippisch O, et al. 2013. Combined impact of soil heterogeneity and vegetation type on the annual water balance at the field scale. *Vadose Zone Journal*, 12(4): 1–17.
- Schwinning S, Sala O E. 2004. Hierarchy of responses to resource pulses in arid and semi–arid ecosystems. *Oecologia*, 141(2): 211–220.
- Shao Q, Gu W, Dai Q Y, et al. 2014. Effectiveness of geotextile mulches for slope restoration in semi-arid northern China. *CATENA*, 116: 1–9.
- Wang S, Fu B J, Gao G Y, et al. 2013. Responses of soil moisture in different land cover types to rainfall events in a re-vegetation catchment area of the Loess Plateau, China. *CATENA*, 101: 122–128.
- Wilson D J, Western A W, Grayson R B. 2004. Identifying and quantifying sources of variability in temporal and spatial soil moisture observations. *Water Resources Research*, 40(2): W02507, doi: 10.1029/2003WR002306.
- Wu H, Li X Y, Li J, et al. 2016. Differential soil moisture pulse uptake by coexisting plants in an alpine *Achnatherum splendens* grassland community. *Environmental Earth Sciences*, 75(10): 914, doi: 10.1007/s12665-016-5694-2.
- Xu G, Zhang T G, Li Z B, et al. 2017. Temporal and spatial characteristics of soil water content in diverse soil layers on land terraces of the Loess Plateau, China. *CATENA*, 158: 20–29.
- Yu T, Feng Q, Si J, et al. 2019. Responses of riparian forests to flood irrigation in the hyper–arid zone of NW China. *Science of the Total Environment*, 648: 1421–1430.
- Yu X, Huang Y, Li E, et al. 2017. Effects of vegetation types on soil water dynamics during vegetation restoration in the Mu Us Sandy Land, northwestern China. *Journal of Arid Land*, 9(2): 188–199.
- Yu X N, Huang Y M, Li E G, et al. 2018. Effects of rainfall and vegetation to soil water input and output processes in the Mu Us Sandy Land, northwest China. *CATENA*, 161: 96–103.
- Zhang X, Zhang L, Zhao J, et al. 2008. Responses of streamflow to changes in climate and land use/cover in the Loess Plateau, China. *Water Resources Research*, 44(7): W00A07, doi: 10.1029/2007WR006711.
- Zhang Z, Si B, Li M, et al. 2020. Deficit and recovery of deep soil water following a full cycle of afforestation and deforestation of apple trees on the Loess Plateau, China. *Water*, 12(4): 989, doi: 10.3390/w12040989.
- Zhao C L, Jia X X, Zhu Y J, et al. 2017. Long–term temporal variations of soil water content under different vegetation types in the Loess Plateau, China. *CATENA*, 158: 55–62.
- Zhou D, Zhao S, Zhu C. 2012. The Grain for Green project induced land cover change in the Loess Plateau: A case study with Ansai County, Shanxi Province, China. *Ecological Indicators*, 23: 88–94.
- Zhu L, Wang Z H, Mao G L, et al. 2014. Water uptake from different soil depths for halophytic shrubs grown in northern area of Ningxia Plain (China) in contrasted water regimes. *Journal of Plant Interactions*, 9(1): 26–34.